

AD-A106 981 MARYLAND UNIV COLLEGE PARK DEPT OF MATHEMATICS  
FINAL SCIENTIFIC REPORT.(U)  
1980 B KEDEM

F/G 12/1

AFOSR-80-0211

UNCLASSIFIED

AFOSR-TR-81-0718

NL

1 10 1  
21 6200 1



END  
DATE  
FILMED  
28  
DTIC

LEVEL

Final Scientific Report, for Subject Grant AFOSR-80-0211

by

Benjamin Kedem  
Principal Investigator  
Department of Mathematics  
University of Maryland  
College Park, Md. 20742

DTIC  
ELECTRIC  
NOV 10 1981

H

AD A106981

In this effort we further studied the implication of the higher order crossings on stationary time series and homogeneous random fields. The Higher Order Crossings Theorem has been extended to homogeneous random fields [2] and the  $\chi^2$  first introduced in the literature in [4] has been further studied. In particular, since the Higher Order Crossings Theorem can be extended to random fields this suggests that  $\chi^2$  can be used in texture discrimination which is indeed the case.

We have studied the power of  $\chi^2$  via simulation techniques of series and fields. As the alternative becomes more distinguishable so is the increase in the power as expected. We still know very little about the eigenvalues of  $\chi^2$  although we have computed some for special cases.

Another aspect we have been studying is graphics in the series analysis [1]. There is no theory of graphics as such, but we were able to prove various graphical results by comparing a series by a random curve. In this connection we mention our work [2] in which the notion of a unit has been advanced in connection with the appearance of binary sequences.

81 11 06 040

Approved for public release;  
distribution unlimited.

AD A106981  
FILE COPY

# 1. THEOREM 1.1. INCREASING RESULTS

1. Let  $\{X(t_1, t_2), t_1, t_2 = 0, \pm 1, \pm 2, \dots\}$  be a zero mean stationary random field. Then if  $U$  is the clipping operator and  $V$  is the difference operator we have

$$\{U_{t_1}^{(k)} V_{t_2}^{(k)} Z(t_1, t_2)\} = \{a, a'\}, \quad k \rightarrow \infty,$$

where

$$a = \begin{matrix} \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \dots 1 & 0 & 1 & 0 & 1 & 0 \dots \\ \dots 0 & 1 & 0 & 1 & 0 & 1 \dots \\ \dots 1 & 0 & 1 & 0 & 1 & 0 \dots \\ \dots 0 & 1 & 0 & 1 & 0 & 1 \dots \\ \dots 1 & 0 & 1 & 0 & 1 & 0 \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{matrix}$$

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
NTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
by	
Distribution/	
Availability Codes	
Avail and/or	
Special	

**A**

and  $a'$  is the same array shifted to the right.

2. Let  $D_{j,N}$  be the total number of symbol changes in the binary array  $\{U_{t_1}^{(j)} V_{t_2}^{(j)} Z(t_1, t_2)\}$ . Then

a)  $\lim_{N \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{D_{j,N}}{N} = 1$

b)  $\lim_{N \rightarrow \infty} \frac{D_{j,N}}{N}$  is almost surely increasing in  $j$ .

3. Let  $\hat{p}_j = D_{j,N}/(N-1)$ ,  $p_j = E\hat{p}_j$ ,  $\hat{p} = (\hat{p}_1, \dots, \hat{p}_{k-1})'$

where  $\Delta_j = D_j - D_{j-1}$  as in [5]. Define

$$P = D_{AG}(p_1, \dots, p_{k-1}) - P P'.$$

Then

$$\psi_N^2 = (N-1)(\hat{p} - p)' P^{-1} (\hat{p} - p)$$

and under regularity conditions

$$\sqrt{N-1} (\hat{p} - p) \rightarrow N(0, V), \quad N \rightarrow \infty.$$

Then

$$\psi_N^2 \rightarrow \sum_{j=1}^{K-1} \delta_j \Sigma_j^2, \quad \Sigma_j \sim N(0,1) \text{ INDEP.}$$

And

$$\bar{v} = \frac{1}{K-1} \sum_{j=1}^K \frac{v_{jj}}{p_j}$$

It follows that in order to study the distribution of  $\psi_N^2$ , a great deal can be learnt from the asymptotic variance of  $\Delta_j / \sqrt{N-1}$ .

4. From many simulations, it can be shown that 28 is a critical value of  $\psi^2$  which roughly corresponds to 0.05 level. Our power calculations follow using this critical value.

TRUE $H_1: \text{ARMA}(2,2)$				FALSE $H_0: \text{AR}(1)$	POWER $p(0.028)$
$\phi_1$	$\phi_2$	$\theta_1$	$\theta_2$	$\phi$	
0.2	0.0	0.8	0.0	0.6	0.993
0.3	0.0	0.6	0.0	0.62	0.920
0.4	0.4	0.0	0.0	0.66	0.874
0.5	0.0	0.5	0.0	0.72	0.806
0.0	0.0	0.6	0.0	0.45	0.583
0.6	0.3	0.0	0.0	0.85	0.540
0.6	0.0	0.3	0.0	0.73	0.370
0.6	0.3	0.5	0.0	0.95	0.464
0.0	0.0	0.5	0.5	0.51	0.139
0.8	0.0	0.0	0.0	0.85	0.024
-0.6	0.0	-0.5	0.0	-0.65	0.104
0.0	0.0	0.3	0.0	0.23	0.080
0.0	0.0	0.0	0.0	0.013	0.046

Similar results hold for random fields with critical value 65.

## II. RESULTS FOR GRAPHICS

Assume  $|Z_t| < A$ , and in  $(-A, A)$  we define a unit process  $\{U_t\}$  independent of  $\{Z_t\}$  and made of independent variates. We have two time series

$Z_1 Z_2 \cdots Z_N$  + Time Series

$U_1 U_2 \cdots U_N$  + "Random Curve"

1. The finite dimensional distributions of a Gaussian process are completely determined by graphical features.
2. The variance of a rounded stationary process can be obtained by counting the number of crossings by  $Z_t$  of the random curve, the number of sojourns of at least two time periods above and below the random curve, and the number of crossings of  $\sqrt{t}Z_t$  of the corresponding random curve.
3. Let  $\{U_t^{(k)}\}$  be the random curve which corresponds to  $\{\sqrt{k}Z_t\}$ . The crossings of the  $U^{(k)}$ -curve by  $\sqrt{k}Z_t$  completely determine the covariance structure of  $\{Z_t\}$ .
4. Let  $Z_t = cZ_{t-1} + U_t$  where  $|c| < 1$  and  $U_t \sim$  Cauchy with characteristic function  $\exp[-(1-|c|)|\lambda|]$ . Then  $Z_t$  is Cauchy with characteristic function  $\exp[-|\lambda|]$ . Note, the moments of  $Z_t$  do not exist. Yet  $c$  is estimable from the axis crossings by  $Z_t$ ,  $t = 1, \dots, N!$

### III. m'th ORDER UNITS FOR BINARY SEQUENCES

An m'th order unit of a binary sequence, is a sub-sequence which starts with 1, ends with m separating 0's and in which each 0-run consists of at most m-1 0's.

In [1] we have various combinatorial results. They lead to:  $I_K(X_t)$  is m'th order Markov and  $S = \sum_{i=1}^N X_i$ ,  $X_i$  being binary, then

$$r(s) \sim \left( \begin{array}{l} \# \text{ of permutations of the} \\ m\text{'th order units with the} \\ \text{free } 0\text{'s} \end{array} \right) \underbrace{P_{10\dots 0}^s}_m \underbrace{P_{00\dots 0}^{n-s-m}}_{m+1}$$

$$\rightarrow e^{-\beta} \beta^s / s!$$

where  $\beta = N P_{100\dots 0}$  and  $P_{10\dots 0} = p(1|0\dots 0)$ ,  $P_{00\dots 0} = p(0|0\dots 0)$ .

Our combinatorial results help to define higher order degeneracies of particles in the physical sense.

## References

- [1] Kedem, E.(1980). Graphical Considerations in Time Series Analysis, University of Maryland Report MD-80-43-BK (Submitted).
- [2] Kedem, E.(1981). On Nearest Neighbor Degeneracies of Indistinguishable Particles, Journal of Math. Physics, 22, 456-461.
- [3] Kedem, E.(1981). A Convergence Phenomenon and Nonparametric Classification of Random Fields. Submitted for publication.
- [4] Kedem, E. and Slud, E. (1981). On Goodness of Fit of Time Series Models, Biometrika, 68, 551-556.
- [5] Kedem, E. and Slud, E. (1979). Higher Order Crossings in the Discrimination of Time Series, I, II, Univ. of Maryland Tech. Reports MD79-73-BK/ES, MD-79-89-BK/ES.



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR. 81 -0718</b>	2. GOVT ACCESSION NO. <b>AD-A106981</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) FINAL SCIENTIFIC REPORT FOR SUBJECT GRANT AFOSP 80-0211		5. TYPE OF REPORT & PERIOD COVERED FINAL, 15 MAR 80 - 31 AUG 81
7. AUTHOR(s) Benjamin Kedem		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Mathematics University of Maryland College Park MD 20742		8. CONTRACT OR GRANT NUMBER(s) AFOSR-80-0211
11. CONTROLLING OFFICE NAME AND ADDRESS Mathematical & Information Sciences Directorate Air Force Office of Scientific Research Bolling AFB DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 1980
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this effort further study was made of the implication of the higher order crossings on stationary time series and homogeneous random fields. The Higher Order Crossings Theorem has been extended to homogeneous random fields and the psi-square first introduced in the literature by the principal investigator has been further studied. In particular, since the Higher Order Crossings Theorem can be extended to random fields this suggests that psi-square can be used in texture discrimination which is indeed the case. The (CONT.)		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ITEM #20, CONT.:

investigators have studied the power of psi-square via simulation techniques of series and fields. As the alternative becomes more distinguishable so is the increase in the power as expected. Another aspect studied is graphics in the series analysis. There is no theory of graphics as such, but the investigators were able to prove various graphical results by clipping a series by a random curve. In this connection the investigators did some work in which the notion of a unit has been advanced in connection with the appearance of binary sequences.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)